



**Manchester
Metropolitan
University**

Hasler-Winter, Samuel (2018) An investigation into the relationship between visual imagery and stress. University of Manchester. (Unpublished)

Downloaded from: <https://e-space.mmu.ac.uk/621612/>

Publisher: University of Manchester

Please cite the published version

<https://e-space.mmu.ac.uk>



An investigation into the relationship between visual imagery and stress

Samuel Hasler-Winter

Supervised by: Nils Muhlert

June 2018

VISUAL IMAGERY AND STRESS

List of Contents

Abstract.....	Page 3
Intro.....	Page 6
Method.....	Page 10
Participants.....	Page 10
Design.....	Page 10
Materials.....	Page 10
Procedure.....	Page 11
Data Analysis.....	Page 12
Results.....	Page 13
Discussion.....	Page 16

List of Figures

Figure 1. Scatter-graph of VVIQ and AUC anxiety scores in women in the stress condition.....	Page 14
Figure 2. Scatter-graph of VVIQ and PSRS scores in women.....	Page 15
Figure 3. Mean VVIQ scores for high and low PSRS groups in women.....	Page 16

List of Tables

Table 1. Number and percentages of participants in the high and low scoring.. groups for PSRS and VVIQ.....	Page 13
Table 2. Number and percentages of participants in the high and low scoring.. groups for PSRS and VVIQ.....	Page 13

VISUAL IMAGERY AND STRESS

Abstract

Visual imagery plays a functional role in memory and emotion processing. Similar links between stress, memory and emotion also exist, leading to the proposed hypothesis that there may also be a functional relationship between stress and visual imagery. In 36 students, trait vividness of visual imagery was assessed in relation to two variables: trait stress reactivity and acute stress response. The Trier Social Stress Test (TSST) and a matched control condition were used to induce stress. The Vividness of Visual Imagery Questionnaire (VVIQ) and Perceived Stress Reactivity Scale (PSRS) were used as trait measures of visual imagery vividness and stress reactivity respectively. Neither the relationship between vividness of visual imagery and stress reactivity or acute stress response were significant. These findings suggest that stress and visual imagery may not be as closely related as thought. However, significant gender differences seen in the sample and proposed in the literature led to exploratory analyses. These analyses revealed a significant effect of stress-reactive group (high/low) on visual imagery in women, such that low stress reactive individuals experienced more vivid visual imagery and vice versa. Additionally, in women only, the hypothesised inverse relationships: between visual imagery vividness and both trait and acute stress were found to have moderate and strong correlations that were approaching significance, respectively. Given that this was post-hoc analysis, firm conclusions cannot be drawn, although this suggests that the proposed inverse relationship between voluntary visual imagery and stress may exist in women in the healthy population.

Key Words: Visual Imagery, Vividness, Stress, Stress-reactivity

VISUAL IMAGERY AND STRESS

Declaration

The contents of this paper are the students original work unless referenced clearly to the contrary. No portion of the work referred to in the dissertation has been submitted in support of an application for another degree or qualification of this or any other university or other institute of learning.

Acknowledgements

Thank you to Liz McManus, Amy Flynn, Nathaniel Crawford and Raphael Dallery for their assistance in the collection of this data, and Nils Muhlert for his supervision.

Dedication

For my parents, Susan and James, for their unwavering love and support. Also, Liz McManus, for keeping me sane despite my best efforts to the contrary.

Intellectual Property Statement

- i. The author of this dissertation (including any appendices and/or schedules to this dissertation) owns certain copyright or related rights in it (the “Copyright”) and s/he has given The University of Manchester certain rights to use such Copyright, including for administrative purposes.
- ii. Copies of this dissertation, either in full or in extracts and whether in hard or electronic copy, may be made only in accordance with the Copyright, Designs and Patents Act 1988 (as amended) and regulations issued under it or, where appropriate, in accordance with licensing agreements which the University has entered into. This page must form part of any such copies made.
- iii. The ownership of certain Copyright, patents, designs, trademarks and other intellectual property (the “Intellectual Property”) and any reproductions of copyright works in the dissertation, for example graphs and tables (“Reproductions”), which may be described in this dissertation, may not be owned by the author and may be owned by third parties. Such Intellectual Property and Reproductions cannot and must not be made available for use without the prior written permission of the owner(s) of the relevant Intellectual Property and/or Reproductions.
- iv. Further information on the conditions under which disclosure, publication and commercialisation of this dissertation, the Copyright and any Intellectual Property and/or Reproductions described in it may take place is available in the University IP Policy, in any relevant Dissertation restriction declarations deposited in the University Library, and The University Library’s regulations.

VISUAL IMAGERY AND STRESS

Introduction

Visual imagery is for many, central to human experience. It has been shown to have a functional role in processes as diverse as memory, regulating emotion, decision-making, and addiction, as well as contributing to the pleasure and persuasiveness of a written narrative (Amit and Greene, 2012; Epstein & Pacini, 2001; Green and Brock, 2002; Kavanagh, Andrade, & May, 2005; Kosslyn et al., 1990; Paivio, 1969; Sparks, Sparks and Gray, 1995).

Visual imagery is commonly understood to be the experience of perceiving an object, scene or person in the mind, in the absence of the relevant external stimuli (de Vito and Bartolomeo, 2016). A phenomena that is comparable to the experience of seeing in typical perception, but often somewhat less defined or vivid (Zeman et al., 2010). Furthering this, Paivio's (1986) dual-coding hypothesis posits that imagery and language are two distinct and additive channels for information processing in the brain, a view upheld by current theorists and supported by evidence of the functional role of visual imagery (Libby and Eibach, 2013). Although there is some debate, visual imagery serves mainly to represent specific, concrete information that is not well represented in language, and doing so serves a function similar to vision (Kosslyn, 2006; Kounios and Holcomb, 1994; Libby and Eibach, 2013; Slee, 1980).

Visual imagination was described by Galton (1880) as a varying trait, suggesting that it's vividness varies between individuals. This suggestion has since been confirmed by neuroscience in more recent years (Pearson, 2014). "Aphantasia", a condition present in 2% of the population involving a complete absence of voluntary mental visual imagery, appears to have both structural and psychogenic causes (de Vito and Bartolomeo, 2016; Zeman, Dewar & Della Sala, 2015; 2016). However, little is known about the cause of variation in visual imagery in the non-clinical population.

Evidence from studies employing a binocular rivalry methodology – in which two differing stimuli are simultaneously shown to the two eyes – demonstrate that the act of perception is altered by imagery priming in much the same way as priming with external stimuli (Pearson, Clifford and Tong, 2008). This effect correlates with subjective reports of vividness in the normal population and in aphantasics (for whom no priming effect is found) (Keogh and Pearson, 2017; Pearson, Rademaker and Tong, 2011). This suggests that in both populations subjectively reported differences in visual imagery vividness reflect genuine differences in phenomenal experience, rather than impaired meta-cognition (Keogh and Pearson, 2017; Pearson et al., 2011). Functional magnetic resonance imaging (fMRI) data shows distinct neural correlates of these subjectively reported and behaviourally measured findings in aphantasics and healthy participants, as well as common neurological underpinnings between visual perception and visual imagery (Cui et al., 2007; Horikawa et al., 2013; Ishai, Haxby and Ungerleider, 2002; Kosslyn et al., 1993; MacKisack et al., 2016; Pearson et al., 2015; Zeman et al., 2010). These data provide strong evidence that visual mental imagery is a visual experience in the phenomenological sense (i.e. resembling perception), and that metacognition and subjective reports of visual imagery are accurate in both healthy and clinical cases. The Vividness of Visual Imagery Questionnaire (VVIQ) is a subjective trait measure

VISUAL IMAGERY AND STRESS

of voluntary visual imagery shown to have high reliability and construct validity, demonstrated partly by meaningful correlation with neural activation (Campos and Pérez-Fabello, 2009; Fulford, 2017; McKelvie, 1995).

Shared cortical activation between visual imagery processing and memory suggest that visual imagery plays a significant role in memory (Aydin, 2018; Farah, 1985). Indeed, a large body of behavioural evidence shows visual imagery plays a functional role in memory, and as such, vivid visual imagery ability, instructions to visualise, or highly imageable and concrete stimuli result in an improved memory performance (Abelson, 1975; Keogh and Pearson, 2011; McKelvie and Demers, 1979; Paivio, 1969; Swann and Miller, 1982; Williams, Healy, & Ellis, 1999). Vividness of visual imagery and memory are also positively correlated in aphasia, severely deficient autobiographical memory, highly superior autobiographical memory and amnesia (Hassabis et al., 2007; LePort et al., 2012; Palombo, 2015; Watkins, 2017; Zeman et al., 2016). In fact, visual imagery plays such a significant role in memory that vividly imagining fictional autobiographical situations can lead to false memories of them actually occurring (Hyman and Pentland, 1996). The relationship between stress and memory is also well documented (Kuhlmann, Piel and Wolf, 2005; Sauro, Jorgensen and Pedlow, 2003). Transient stress influences both encoding and retrieval memory processes (Domes et al., 2002; Roozendaal, 2002; Schwabe and Wolf, 2010). Although the direction of the effect depends on various factors, crucially, stress can have both enhancing and damaging effects on memory (Cahill, Gorski and Le, 2003; Diamond et al., 2007; Schwabe et al., 2012). In clinical cases, extreme stress can cause amnesia (Freyd, 1994), or memory flashbacks in post-traumatic stress disorder (PTSD) (Ehlers, Hackmann & Michael, 2004). The functional involvement of visual imagery in memory, and the strong influence of stress on memory, taken together, may suggest a link between stress and visual imagery.

The process by which the body responds to stress is referred to as allostasis (McEwen, 2000; 2005). Allostasis is a process of adaptation managed mainly by the Autonomic Nervous System and the Hypothalamic-Pituitary-Adrenal (HPA) axis (McEwen, 1998). This biological reactivity to stressors is evolutionarily selected to prepare the individual for a threat (Ellis, Jackson and Boyce, 2006). However, an excess of these hormones can contribute to or exacerbate disease (McEwen and Seeman, 1999). In times of severe stress, the HPA axis produces excess cortisol, which has a dysregulatory effect on the HPA axis, as well as influencing a number of brain areas including the hippocampus, a core brain region for memory-related processes, including visual imagery (Fortin, Agster and Eichenbaum, 2002; Ishai et al., 2002; Leonard and Myint, 2009; Miller and O'Callaghan, 2002; Sapolsky, 1996). This maladaptation, referred to as allostatic load, describes the dysregulation of stress responses and is one of a variety of factors that can influence how an individual will respond to stress (McEwen and Stellar, 1993). In doing so, the presence of allostatic load results in the negative contributions of stress, including a number of clinical conditions (Dallman, 2010; Seeman et al., 1997; Selye, 1955; Steptoe and Kivimäki, 2012).

Especially powerful in determining the presence of allostatic load are developmental experiences, which impact lifelong physiological reactivity to stress due to increased

VISUAL IMAGERY AND STRESS

susceptibility to environmental influences during this time (Heim and Binder, 2012; Hertzman and Boyce, 2010). Referred to as epigenetics, the interaction between these environmental stressors and genes, particularly at these sensitive points, is responsible for many of the individual differences we see in reactivity to daily stressors (Provençal and Binder, 2015). Evidence shows that individuals who experience severe early life trauma display significantly increased stress reactivity (Flinn and England, 1995; Heim et al., 2000). Psychological processes also mediate the impact of stressful life events, although the formation of these themselves is dependent on early life stress (Hankin, 2005; Young, Klosko and Weishaar, 2003). Other contributing factors are physiological vulnerability: resulting from both genetic predisposition and elements such as diet, exercise, substance use and gender (Claessens et al., 2011; McEwen, 2006; Verma et al., 2011).

Although all forms of stress elicit similar HPA activation, distinctions must be made between physical and social stressors as these impact upon different neural circuits and therefore may have different cognitive effects (Dedovic et al., 2009). Social Self Preservation Theory states that situations which threaten the 'social-self' by involving a social-evaluative component, pose a risk to an individual's self-esteem and in doing so activate the HPA axis (Gruenewald et al., 2004). One method of measuring these individual differences in social stress reactivity is the Perceived Stress Reactivity Scale (PSRS), a subjective tool found to have good stability over time and convergent validity with related constructs (Schlotz et al., 2011).

Longitudinal studies of Cushing's syndrome, which features spontaneous hypercortisolism, show that this is followed in many cases by emotional disturbances (Starkman et al., 1981). This highlights the potential that heightened cortisol levels contribute to dysfunction in emotion regulation and helps imply the direction of causation in an already acknowledged association between cortisol, stress and emotion dysregulation (El-Sheikh et al., 2008; Keenan, 2000). In contrast, visual imagery has been shown to aid interpretation and regulation of emotions, suggesting that visual imagery may in some way mediate this relationship between stress and emotion processing (Dirkx, 2001; Kosslyn et al., 1990). Emotional suppression is also associated with decreased vividness of imagery for both imagined and remembered events, as well as greater stress generally (D'Argembeau and der Linden, 2006; Iwamitsu et al., 2009; Moore, Zoellner & Mollenholt, 2008). This suggests stress may be one mechanism by which visual imagery vividness is inhibited, or that deficits in visual imagery vividness contribute to heightened stress reactivity, or indeed, that the relationship is bi-directional. An alternative explanation is that detrimental individual differences in stress reactivity and visual imagery vividness are both the result of similar maladaptations in the HPA axis although are not functionally related. However, a functional relationship between stress and visual imagery better accounts for the effectiveness of imagery use in therapies for PTSD, social phobia, OCD, depression and a range of other anxious disorders - all disorders characterised by extreme stress reactivity and emotion dysregulation (Cisler and Olatunji, 2012; Etkin and Wagner, 2007; Foa et al., 1980; Hirsch and Holmes, 2007; Holmes, Arntz and Smucker, 2007; Krakow, Hollifield and Johnston, 2001; Speckens et al., 2007).

VISUAL IMAGERY AND STRESS

There are also a number of clinical differences in visual imagery. A key distinction in these differences is between voluntary and involuntary (or intrusive) imagery. Voluntary imagery refers to imagery experienced due to a conscious (although not necessarily effortful) act of intent to picture something, and involuntary imagery being experienced spontaneously without any such intent (Homer and Deeprose, 2017). In clinical cases characterised by involuntary imagery, an excess of this type of imagery is often found (Brewin et al., 2010; Ehlers et al., 2004; Speckens et al., 2007). In contrast, disorders featuring decreases in visual imagery primarily concern voluntary visual imagery (Sierra, 2009). This dissociation is supported by the selective absence of voluntary visual imagery in aphantasia, with the preservation of involuntary imagery (Zeman et al., 2015).

As mentioned previously, PTSD can cause hyper-activation and vivid, involuntary re-experiencing of memories (Ehlers et al., 2004). Eating disorders, which are often greatly exacerbated by stress, also feature highly vivid and distressing involuntary visual imagery (Brewin et al., 2010). Intrusive visual imagery is also sometimes found in Obsessive Compulsive Disorder (OCD), with those who experience visual imagery having more obsessive beliefs, compulsive behaviours and anxiety than those who do not (Speckens et al., 2007). OCD sufferers also exhibit significantly greater cortisol and perceived stress relative to healthy individuals and their perceived stress correlates with the severity of their condition (Morgado et al., 2013). This strongly reinforces an association between stress, cortisol, and visual imagery, and also suggests that psychogenic deletion of visual imagery in such disorders may serve an adaptive function.

In contrast, stress-related disorders such as depression, anxiety and depersonalization are all associated with diminished experience of voluntary, and in particular positive, visual imagery (Holmes et al., 2016; Sierra, 2009). In the context of depression, Paivio's (1985) imagery framework is highly relevant, suggesting that imagery plays a function in managing motivation, arousal and affect, as well as representing goals and activities. Of course, depression is characterised not only by a lack of motivation, but dysregulation of arousal and affect, as well as a loss of interest and pleasure in goals and activities (American Psychiatric Association, 2013). This framework therefore outlines several mechanisms by which a deficit in visual imagery vividness could contribute to the excessive stress characteristic of depression.

Taking the clinical data as a whole, it suggests that stress is related to both hypoactive voluntary visual imagery and hyperactive involuntary visual imagery. Of course, clinical samples are biased towards more extreme levels of stress and visual imagery, and therefore associations are more likely to be found. However, the maladaptations of stress response that underlie many clinical disorders do not only exist within these clinical parameters suggesting that these clinical relationships between individual differences in visual imagery and stress may be mirrored in more subtle ways amongst healthy individuals (Claessens et al., 2011). Additionally, the non-clinical data also suggests that the relationship between stress and voluntary visual imagery is negatively correlated. There is also limited evidence to suggest there is a relationship between stress and hyperactive involuntary imagery in the non-clinical population (Berntsen and Rubin, 2008; Witvliet, 1997). However,

VISUAL IMAGERY AND STRESS

presently there is no empirical research directly investigating the relationship between stress and visual imagery (voluntary or involuntary) within the normal population. Although the topic of involuntary visual imagery requires further understanding, the focus of this investigation is voluntary visual imagery and as such, it is expected that an inverse relationship between stress and visual imagery will be found, if any.

One method of manipulating stress in the non-clinical population is the Trier Social Stress Test (TSST). This task uses a social-evaluative and uncontrollable paradigm to replicate the influence of life's daily social stressors and reliably induce large cortisol responses in healthy participants (Kirschbaum, Pirke and Hellhammer, 1993). Compared to other forms of social stress, some using just public speaking and others using different paradigms, the Trier elicits greater and more robust cortisol elevations making it the obvious choice of task (Dickerson and Kemeny, 2004). Measuring self-report anxiety and salivary cortisol response during this task, coupled with later measures of trait stress reactivity and trait visual imagery, this study aims to explore the relationship between trait visual imagery, trait stress reactivity and acute stress response in a population of undergraduate students. The primary hypothesis of this study predicts that there will be a negative relationship between trait vividness of visual imagery and trait stress reactivity. A secondary hypothesis predicts that there will be a negative relationship between trait vividness of visual imagery and acute stress response.

Method

Participants

39 participants were opportunistically recruited from a pool of undergraduate psychology students at The University of Manchester (UoM). 3 participants withdrew from the study. The age of the remaining 36 participants (24 female, 12 male) ranged from 18-26 years old ($M = 20.66$, $SD = 2.179$). Participants were split randomly between two conditions (18 stress, 18 control). Informed consent was received from all participants. Participants were obtained from a non-clinical population and therefore were not at any identified risk from stress induction. Participants were awarded university credits or cash payment for their participation. All data were de-identified to ensure confidentiality. An independent ethics committee approved the experiment.

Design

The experiment used a between-subjects design, with participants randomly allocated between two conditions (18 stress, 18 control). The experimental design was also correlational. Independent variables were condition, voluntary visual imagery vividness and perceived stress reactivity. The dependent variable was acute stress response.

Materials

VISUAL IMAGERY AND STRESS

A modified version of the Trier Social Stress Test (TSST) (Kirschbaum et al., 1993) was used to induce stress. The Vividness of Visual Imagery Questionnaire (VVIQ) (Marks, 1995) was used to subjectively measure trait voluntary visual imagery (see Appendix A). Trait stress reactivity was measured using the Perceived Stress Reactivity Scale (PSRS) (Schlotz et al., 2011) (see Appendix B). Salivary cortisol was measured as a biomarker of psychological stress response (Choi et al., 2014) – this was analysed through ELISA kits, a precise, reliable tool (Shimada et al., 1995). Stress response was also assessed using a Visual Analogue Scale for Anxiety (VAS-A) - shown to be a reliable, valid, sensitive measure with good convergent validity (Cella and Perry, 1976; Facco et al., 2011; 2013; Hornblow and Kidson, 1976) (see Appendix C).

Procedure

Testing was carried out in the afternoon for all participants in order to control for the influences of circadian rhythm on resting cortisol levels (Thuma et al., 1995). Participants were required to abstain from eating and drinking 2-hours prior to arrival. Upon arrival, participants were given an information sheet about the study (see Appendix D), informing them they would be involved in a stress test. They were then required to sign consent forms (see Appendix E). Following this, there was a ten-minute waiting period to allow participant cortisol to return to resting levels. Participants were then questioned briefly regarding that day's smoking habits, exercise, illness and medication (see Appendix F for list of questions), as these factors are all known to influence cortisol levels (Direk et al., 2011; Hibel et al., 2006; Hill et al., 2008; Kirschbaum, Pirke and Hellhammer, 1995). Participants then completed a Visual Analogue Scale for Anxiety (VAS-A) and provided a saliva sample.

Participants in the stress condition were then taken into a second testing room to be introduced to the TSST, where they were introduced to the panel and told they were required to give a 5-minute speech for a job interview for a position at the UoM Psychology department, after which another task would follow. They were informed that the interview was video and voice recorded for analysis and that one panel member was trained in behavioural analysis and was taking notes (see Appendix G for experimenter script). Participants in the control condition were taken into the second testing room and introduced instead to the control task, which required them to speak about a topic of their choice for 5 minutes with friendly peers. In this condition, normal positive social cues from the panel were permitted although the participant was still encouraged to lead the discussion. Following introduction to the stress and control tasks respectively participants were returned to the initial testing room for a ten-minute preparation period where they were allowed to make notes.

After the 10-minute waiting period participants completed a second VAS-A. Control and stress condition participants then returned to the second testing room to have their discussion or give their speech respectively. The stress-condition speech was given unaided (without notes) to a panel of two unresponsive "job interviewers" who offered no positive social cues to the participant. Periods of silence were allowed for 20 seconds before a follow up question was asked (see Appendix H for interviewer script) After the speech/discussion participants completed a third VAS-A. Following

VISUAL IMAGERY AND STRESS

this, participants were instructed regarding and subsequently completed the second task. For the stress condition this was a 5-minute mental arithmetic task that involved counting backwards out loud in 13s from 1687. For the control maths task a booklet of simple addition and subtraction questions was completed by the participant (see Appendix I). Participants then completed a fourth VAS-A and provided a second saliva sample. Following this, both stress and control condition participants were returned to the initial testing room for a 10-minute resting period, after which they completed a final VAS-A and saliva sample. Participants in the stress condition were then debriefed regarding the nature of the stress test, that the job interview and behavioural analyst were fictitious, that the video camera and voice recorder were not functional and that the panel were deliberately unresponsive and abstaining from positive feedback with the purpose of provoking a stress response. Participants returned the following day, signed a second consent form and completed the VVIQ and the PSRS.

Data Analysis

The area under the curve for VAS-A (AUC anxiety) scores was found to estimate each individual's change in anxiety during the TSST or control task. An independent samples t-test was carried out between AUC anxiety for stress and control conditions to test the effectiveness of our modified TSST. A Pearson's correlation co-efficient on PSRS and AUC anxiety scores was used to assess the convergent validity of these two measures in the sample.

A median split was performed on VVIQ scores to form high and low VVIQ groups. PSRS scores were then tested for measures of normality, these were met and so analysis proceeded with parametric tests. An independent samples t-test was conducted with an independent variable of VVIQ group and dependent variable of PSRS. Pearson's correlation co-efficient was also conducted between PSRS and VVIQ scores.

For analysis involving AUC anxiety and cortisol-response scores data were split between stress and control conditions and stress data only used. AUC anxiety (stress) scores were tested for measures of normality. Measures of normality were met so analysis proceeded with parametric tests. An independent samples t-test was performed with an independent variable of VVIQ group (stress) and dependent variable of AUC anxiety (stress). Pearson's correlation co-efficient was then carried out between AUC anxiety scores (stress) and VVIQ scores (stress), and between cortisol-response (stress) and VVIQ scores (stress).

Normality of data was tested in each case by calculating the z-score for skewness ($\text{skewness}/\text{SEskewness}$) and kurtosis ($\text{excess kurtosis}/\text{SEexcess kurtosis}$), for a sample size < 50 , z-scores < 1.96 were judged to be normal (Kim, 2013).

Due to the observed gender differences between conditions, exploratory analysis were carried out to assess the role of gender differences, if any.

Results

The independent samples t-test of AUC anxiety revealed there was a significant difference in the scores for stress ($M=21.60$, $SD=4.81$, $SE=1.13$) and control ($M=12.68$, $SD=7.55$, $SE=1.78$) conditions; $t(34)=3.76$, $p=.001$, $SE=2.11$ and 95% CI [3.63, 12.21]. Table 1 shows descriptive statistics of number and age of participants in each gender between conditions. Pearson's correlation co-efficient of AUC anxiety and PSRS scores returned a significant moderate correlation between the two variables ($r=.376$, $n=30$, $p=.026$).

	Stress			Control		
	Females	Males	Total	Females	Males	Total
N	11	7	18	13	5	18
Age	20.50	20.57	20.53	20.73	21.00	20.80
	(2.07)	(2.23)	(2.07)	(2.45)	(2.45)	(2.37)

[illegible]

An independent samples t-test revealed there was not a significant difference in the PSRS scores for High VVIQ ($M = 23.83$, $SD = 7.55$, $SE = 1.78$) and Low VVIQ ($M = 23.47$, $SD = 7.71$, $SE = 1.87$) groups; $t(33) = -.141$, $p = .731$. PSRS was normally distributed, with skewness of $-.221$ ($SE = 3.98$) and kurtosis of $-.168$ ($SE = .778$). Pearson's correlation co-efficient between PSRS and VVIQ was not significant ($r = -.163$, $n = 35$, $p = .349$).

	□□□□□□□□				□□□□□□□□			
	□□□□□□		□□□□		□□□□□□		□□□□	
□□□□□□	□□□□□□□□		□□□□□□		□□□□□□□□		□□□□□□□□	
□□	□□		□□□		□□□		□	
□□□□□□□	□□□□□□□□		□□□□□□		□□□□□□□□		□□□□□□□□	
□□	□□		□□□		□□		□□	



Table 2. shows the number and percentages of male and female participants in the high and low groups for VVIQ and PSRS scores. There was a disproportionately large number of women in the high scoring participant groups for both PSRS and VVIQ measures.

VISUAL IMAGERY AND STRESS

An independent samples t-test revealed there was no significant difference in the AUC anxiety scores between High VVIQ ($M = 19.94$, $SD = 2.55$, $SE = .96$) and Low VVIQ ($M = 21.02$, $SD = 5.90$, $SE = 1.78$) conditions; $t(15) = -.535$, $p = .601$. Levene's test for equality of variances was significant, $F = 6.671$, $p = .020$. Owing to this violation, a t statistic not assuming homogeneity of variance was calculated. AUC anxiety was normally distributed, with skewness of .022 ($SE = .393$) and kurtosis of -.363 ($SE = .768$). Pearson's correlation co-efficient between AUC anxiety scores (stress) and VVIQ scores (stress) showed no significant correlation between the two variables ($r = -.280$, $n = 18$, $p = .260$).

Exploratory analyses were carried out to examine gender differences and findings within each gender separately. An independent samples t-test revealed there was no significant difference between VVIQ scores between men ($M = 52.08$, $SD = 17.59$, $SE = 5.08$) and women ($M = 56.92$, $SD = 10.44$, $SE = 2.13$); $t(34) = -1.037$, $p = .307$. An independent samples t-test of AUC anxiety (stress condition) revealed there was a significant difference between men ($M = 17.76$, $SD = 2.60$, $SE = .98$) and women ($M = 22.40$, $SD = 5.10$, $SE = 1.54$); $t(16) = -2.212$, $p = .042$. An independent samples t-test revealed there was a significant difference in PSRS scores between men ($M = 18.08$, $SD = 7.22$, $SE = 2.08$) and women ($M = 26.57$, $SD = 5.95$, $SE = 1.24$); $t(33) = -3.72$, $p = .001$.

A pearson's correlation between VVIQ (stress) and AUC anxiety (stress) in women revealed a large negative correlation that was approaching significance ($r = -.516$, $n = 13$, $p = .071$). The results of this correlation can be found in Figure 1.




Figure 1. Scatter-graph of VVIQ and AUC anxiety scores in women in the stress condition.

VISUAL IMAGERY AND STRESS

A Pearson's correlation between VVIQ and PSRS scores in women revealed a moderate negative correlation that was approaching significance ($r = -.389$, $n = 23$, $p = .060$). The results of this correlation can be found in Figure 2.




Figure 2. Scatter-graph of VVIQ and PSRS scores in women.

An independent samples t-test was performed on PSRS scores (women) between high VVIQ (women) and low VVIQ (women) groups. Levene's test for equality of variances was significant, $F = 4.395$, $p = .048$. Owing to this violation, a t statistic not assuming homogeneity of variance was calculated. The t-test showed no significant difference in PSRS scores for low VVIQ ($M = 27.78$, $SD = 4.44$, $SE = 1.48$) and high VVIQ ($M = 25.79$, $SD = 6.80$, $SE = 1.82$) groups in women; $t(20.97) = .850$, $p = .405$. A median split was performed on PSRS scores to create a high PSRS and low PSRS group. An independent samples t-test revealed that there was a significant difference between VVIQ scores for high PSRS ($M = 53.44$, $SD = 10.44$, $SE = 2.61$) and low PSRS ($M = 63.88$, $SD = 6.45$, $SE = 2.23$) groups in women; $t(22) = 2.577$, $p = .017$. Figure 3 shows mean VVIQ scores for high and low PSRS groups in women.

VISUAL IMAGERY AND STRESS




Figure 3. Mean VVIQ scores for high PSRS and low PSRS groups in women.

Discussion

Analyses of the relationship between trait stress reactivity and trait visual imagery were not significant. This finding suggests that there is no relationship between trait stress reactivity and trait voluntary visual imagery. Likewise, tests of the relationship between acute stress response and trait vividness of visual imagery were not significant. This suggests that there is also no relationship between an individual's stress response to an environmental stressor and their trait vividness of visual imagery. The independent samples t-test performed on self-report anxiety between conditions revealed a significant effect of the modified TSST in inducing a stress response in participants. This demonstrates that the lack of a relationship found between acute stress and visual imagery was not the result of inadequate stress induction by our task. The significant moderate correlation between measures of acute stress response and trait stress reactivity attests to the convergent validity of these measures in our sample.

The lack of relationship observed between visual imagery and both acute and trait stress is contrary to the proposed hypothesis devised based on previous research linking both stress and visual imagery to memory and emotion regulation (Ehlers et al., 2004; Foa et al., 1980; Freyd, 1994; Hassabis et al., 2007; Sierra, 2009; Starkman et al., 1981; Zeman et al., 2016). This may be in part due to the inability to apply clinical findings to healthy populations. However, the same rationale can also be made based on non-clinical data (Aydin, 2018; Dirkx, 2001; D'Argembeau and der Linden, 2006; Keogh and Pearson, 2011; Kosslyn et al., 1990; Moore et al.,

VISUAL IMAGERY AND STRESS

2008; Schwabe and Wolf, 2012). Sample size was determined by a power calculation to ensure it was sufficiently powered to detect a moderate effect, so inadequate power cannot be responsible for the finding of no relationship in the sample. Practical limitations meant that cortisol data were unable to be analysed in time for submission of this report, however, cortisol data would be expected to correlate strongly with self-report acute anxiety as obtained by the VAS-A scale (Saliba et al., 2016). As such, the absence of this data does not pose a serious limitation to this study. However, it is possible that gender differences may have obscured any effects between visual imagery and stress. With respect to gender differences, cortisol measurements may have been informative, given that research shows that men exhibit significantly larger salivary cortisol measurements following stress exposure in the TSST (Kudielka et al., 2007).

Gender differences were significant in the case of stress reactivity and acute stress response, with women scoring higher in each case. However, no significant gender differences were observed in visual imagery vividness. These gender differences led to a separate analysis of the relationship between stress and visual imagery in women only (men were a very small part of the sample, and revealed no interesting or significant findings). A significant difference in vividness of visual imagery was found between high and low stress reactive groups. This suggests that the hypothesised inverse relationship between voluntary vividness of visual imagery and stress reactivity may exist in women. The negative correlation between vividness of visual imagery and perceived stress reactivity in women was approaching significance, as was the negative correlation between vividness of visual imagery and acute stress response to the TSST in women, supporting this interpretation. Due to the smaller sample size created by splitting data based on gender (and by gender and condition for analysis involving acute stress response), these exploratory analyses are underpowered. Therefore, further investigation in women with a larger sample size is required to confirm whether or not these relationships would each reach significance respectively. However, given that stress response and stress reactivity are related constructs, the approaching significant correlation between both of them and visual imagery, along with the significant difference in visual imagery vividness found between high and low stress reactive groups, implies that this may not be a chance statistical occurrence – and that stress and vividness of voluntary visual imagery may indeed be inversely related in women. No significant difference was found in stress reactivity between high and low vividness groups. This suggests that whereas high or low stress reactivity can be used as a predictor of visual imagery vividness, visual imagery vividness cannot be used as a predictor of stress reactivity.

The finding that women have both a higher acute stress response and greater stress reactivity than men is supported by the consensus in the literature that women have a higher self-report anxiety than men (Milani et al., 2004; Piccinelli and Wilkinson, 2000; Schlotz et al., 2011). In contrast, previous findings show that women also exhibit more vivid visual imagery, however our data found no significant difference between genders in visual imagery (Campos and Sueiro, 1993; Richardson, 1995). This may be the result of these analyses being underpowered due to the gender split rather than a genuine absence of effect. The finding that women's VVIQ scores were

VISUAL IMAGERY AND STRESS

on average higher than men's supports this interpretation. These analyses generally support the understanding that our data is representative of the wider population.

Future research could investigate the relationship between stress and vividness of visual imagery in women specifically, with a larger sample, to enable firm conclusions to be drawn about the relationship between these two variables in this population. Investigation into these effects in men is also required but crucially this must be done separately, due to clear gender differences. Future research could also employ a state measure of visual imagery vividness such as a trial-by-trial subjective report as this has been shown to have significantly larger effect size estimates in correlations with neurological and behavioural measures than the VVIQ (Runge et al., 2017). A trial-by-trial methodology would also be capable of identifying transient changes in visual imagery vividness due to stress and help to infer causality. Alternatively, behavioural measures of imagery such as binocular rivalry would offer the benefit of not being susceptible to socially desirable responding, as the VVIQ has been shown to be (Allbutt et al., 2008; 2011). It is widely acknowledged in the literature that visual imagery is not an entirely unified construct and can be split into object and spatial imagery (Aydin, 2018; Blajenkova, Kozhevnikov and Motes, 2006a; 2006b; Vannucci and Mazzoni, 2009). Another potential direction for future research in this field may then be for the analysis of the relationship between stress and each of these distinct subtypes of visual imagery.

This investigation is not adequate to make such claims, but if it were found that low visual imagery vividness is reliably associated with heightened stress reactivity, then low visual imagery vividness could represent a risk factor for stress-related disorders such as depression or anxiety. Indeed, one potential method for reducing stress in individuals and helping them avoid entering clinical cycles of stress could be the training of visual imagery to enhance vividness. One study attempted to train increased vividness of visual imagery and failed, however this study had only a 5-day time frame, which could be a serious limitation (Rademaker and Pearson, 2012). A study by Aleman et al. (2000) found that musically trained participants had superior auditory imagery ability for both musical and non-musical auditory imagery. Whilst evidence suggests these effects are not cross-modal, it does seem likely that if training of imagery is possible in one modality it could be in others (Aleman et al., 2000). However, this study was correlational and therefore cannot infer causality. It is possible that people with superior auditory imagery ability are more inclined to acquire musical training, for instance. Future research could investigate this question in the context of visual imagery; perhaps with photographic or visual art training. Blackwell et al. (2015) found that in depressives, practising generating positive imagery in response to ambiguous scenarios lead to an increase in positive involuntary imagery experienced in daily life, with beneficial behavioural outcomes and reduction in anhedonic symptoms of depression. Crucially, imagery vividness during the first session of the intervention was a strong predictor of the degree in ultimate reduction of depressive symptoms (Blackwell et al., 2015). Therefore, if a viable method of training increased vividness of visual imagery was achieved, this could be used in conjunction with a mode of therapy such as this to significantly enhance efficacy.

VISUAL IMAGERY AND STRESS

In conclusion, this research contributes to the study of visual imagery by being the first of its kind as a direct investigation into the relationship between visual imagery and stress in a healthy population. In addition, this study helps to clarify that research in this area should be gender specific. This is because there are significant differences between genders with regards to trait stress, acute stress and trait vividness of visual imagery, which may obscure effects if left uncontrolled for. This study demonstrated that no relationship between either acute or trait stress and vividness of visual imagery is found unless controlling for gender. Controlling for gender, however, indicates a possible negative relationship of vividness of visual imagery with both acute stress and trait stress reactivity in women only. This may have valuable theoretical and therapeutic implications for clinical psychology and the study of psychological wellbeing, as well as the continued understanding of both visual imagery and stress fields in their own right.

VISUAL IMAGERY AND STRESS

References

- Abelson, R. P. (1975, June). Does a story understander need a point of view?. In *Proceedings of the 1975 workshop on Theoretical issues in natural language processing* (pp. 140-143). Association for Computational Linguistics.
- Aleman, A., Nieuwenstein, M. R., Böcker, K. B., & de Haan, E. H. (2000). Music training and mental imagery ability. *Neuropsychologia*, 38(12), 1664-1668.
- Allbutt, J., Ling, J., Heffernan, T. M., & Shafiullah, M. (2008). Self-report imagery questionnaire scores and subtypes of social-desirable responding: Auditory imagery, visual imagery, and thinking style. *Journal of Individual Differences*, 29(4), 181-188.
- Allbut, J., Ling, J., Rowley, M., & Shafiullah, M. (2011). Vividness of visual imagery and social desirable responding: correlations of the vividness of visual imagery questionnaire with the balanced inventory of desirable responding and the Marlowe–Crowne scale. *Behavior research methods*, 43(3), 791.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Amit, E., & Greene, J. D. (2012). You see, the ends don't justify the means: Visual imagery and moral judgment. *Psychological science*, 23(8), 861-868.
- Aydin, C. (2018). The differential contributions of visual imagery constructs on autobiographical thinking. *Memory*, 26(2), 189-200.
- Berntsen, D., & Rubin, D. C. (2008). The reappearance hypothesis revisited: Recurrent involuntary memories after traumatic events and in everyday life. *Memory & cognition*, 36(2), 449-460.
- Blackwell, S. E., Browning, M., Mathews, A., Pictet, A., Welch, J., Davies, J., ... & Holmes, E. A. (2015). Positive imagery-based cognitive bias modification as a web-based treatment tool for depressed adults: a randomized controlled trial. *Clinical Psychological Science*, 3(1), 91-111.
- Blajenkova, O., Kozhevnikov, M., & Motes, M. A. (2006). Object-spatial imagery: a new self-report imagery questionnaire. *Applied Cognitive Psychology*, 20(2), 239-263.
- Blajenkova, O., Kozhevnikov, M., & Motes, M. A. (2006). Object and spatial imagery: distinctions between members of different professions. *Cognitive Processing*, 7(1), 20-21.
- Brewin, C. R., Gregory, J. D., Lipton, M., & Burgess, N. (2010). Intrusive images in psychological disorders: characteristics, neural mechanisms, and treatment implications. *Psychological review*, 117(1), 210.

VISUAL IMAGERY AND STRESS

Cahill, L., Gorski, L., & Le, K. (2003). Enhanced human memory consolidation with post-learning stress: interaction with the degree of arousal at encoding. *Learning & memory*, 10(4), 270-274.

Campos, A., & Pérez-Fabello, M. J. (2009). Psychometric quality of a revised version Vividness of Visual Imagery Questionnaire. *Perceptual and Motor Skills*, 108(3), 798-802.

Campos, A., & Sueiro, E. (1993). Sex and age differences in visual imagery vividness. *Journal of Mental Imagery*.

Cella, D. F., & Perry, S. W. (1986). Reliability and concurrent validity of three visual-analogue mood scales. *Psychological reports*, 59(2), 827-833.

Choi, S., Kim, S., Yang, J. S., Lee, J. H., Joo, C., & Jung, H. I. (2014). Real-time measurement of human salivary cortisol for the assessment of psychological stress using a smartphone. *Sensing and Bio-Sensing Research*, 2, 8-11.

Cisler, J. M., & Olatunji, B. O. (2012). Emotion regulation and anxiety disorders. *Current psychiatry reports*, 14(3), 182-187.

Claessens, S. E., Daskalakis, N. P., van der Veen, R., Oitzl, M. S., de Kloet, E. R., & Champagne, D. L. (2011). Development of individual differences in stress responsiveness: an overview of factors mediating the outcome of early life experiences. *Psychopharmacology*, 214(1), 141-154.

Cui, X., Jeter, C. B., Yang, D., Montague, P. R., & Eagleman, D. M. (2007). Vividness of mental imagery: individual variability can be measured objectively. *Vision research*, 47(4), 474-478.

Dallman, M. F. (2010). Stress-induced obesity and the emotional nervous system. *Trends in Endocrinology & Metabolism*, 21(3), 159-165.

D'Argembeau, A., & Van der Linden, M. (2006). Individual differences in the phenomenology of mental time travel: The effect of vivid visual imagery and emotion regulation strategies. *Consciousness and cognition*, 15(2), 342-350.

Dedovic, K., Duchesne, A., Andrews, J., Engert, V., & Pruessner, J. C. (2009). The brain and the stress axis: the neural correlates of cortisol regulation in response to stress. *Neuroimage*, 47(3), 864-871.

de Vito, S., & Bartolomeo, P. (2016). Refusing to imagine? On the possibility of psychogenic aphantasia. A commentary on Zeman et al.(2015). *Cortex; a journal devoted to the study of the nervous system and behavior*, 74, 334.

Diamond, D. M., Campbell, A. M., Park, C. R., Halonen, J., & Zoladz, P. R. (2007). The temporal dynamics model of emotional memory processing: a synthesis on the neurobiological basis of stress-induced amnesia, flashbulb and traumatic memories, and the Yerkes-Dodson law. *Neural plasticity*, 2007.

VISUAL IMAGERY AND STRESS

Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychological bulletin*, 130(3), 355.

Direk, N., Newson, R. S., Hofman, A., Kirschbaum, C., & Tiemeier, H. (2011). Short and long-term effects of smoking on cortisol in older adults. *International Journal of Psychophysiology*, 80(2), 157-160.

Dirkx, J. M. (2001). The power of feelings: Emotion, imagination, and the construction of meaning in adult learning. *New directions for adult and continuing education*, 2001(89), 63-72.

Domes, G., Heinrichs, M., Reichwald, U., & Hautzinger, M. (2002). Hypothalamic–pituitary–adrenal axis reactivity to psychological stress and memory in middle-aged women: High responders exhibit enhanced declarative memory performance. *Psychoneuroendocrinology*, 27(7), 843-853.

Ehlers, A., Hackmann, A., & Michael, T. (2004). Intrusive re-experiencing in post-traumatic stress disorder: Phenomenology, theory, and therapy. *Memory*, 12(4), 403-415.

Ellis, B. J., Jackson, J. J., & Boyce, W. T. (2006). The stress response systems: Universality and adaptive individual differences. *Developmental Review*, 26(2), 175-212.

El-Sheikh, M., Erath, S. A., Buckhalt, J. A., Granger, D. A., & Mize, J. (2008). Cortisol and children's adjustment: The moderating role of sympathetic nervous system activity. *Journal of Abnormal Child Psychology*, 36(4), 601-611.

Epstein, S., & Pacini, R. (2001). The influence of visualization on intuitive and analytical information processing. *Imagination, Cognition and Personality*, 20(3), 195-216.

Etkin, A., & Wager, T. D. (2007). Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164(10), 1476-1488.

Facco, E., Stellini, E., Bacci, C., Manani, G., Pavan, C., Cavallin, F., & Zanette, G. (2013). Validation of visual analogue scale for anxiety (VAS-A) in preanesthesia evaluation. *Minerva anestesologica*, 79(12), 1389-1395.

Facco, E., Zanette, G., Favero, L., Bacci, C., Sivoletta, S., Cavallin, F., & Manani, G. (2011). Toward the validation of visual analogue scale for anxiety. *Anesthesia progress*, 58(1), 8-13.

VISUAL IMAGERY AND STRESS

Farah, M. J. (1985). Psychophysical evidence for a shared representational medium for mental images and percepts. *Journal of Experimental Psychology: General*, 114(1), 91.

Flinn, M. V., & England, B. G. (1995). Childhood stress and family environment. *Current Anthropology*, 36(5), 854-866.

Foa, E. B., Steketee, G., Turner, R. M., & Fischer, S. C. (1980). Effects of imaginal exposure to feared disasters in obsessive-compulsive checkers. *Behaviour Research and Therapy*, 18(5), 449-455.

Fortin, N. J., Agster, K. L., & Eichenbaum, H. B. (2002). Critical role of the hippocampus in memory for sequences of events. *Nature neuroscience*, 5(5), 458.

Freyd, J. J. (1994). Betrayal trauma: Traumatic amnesia as an adaptive response to childhood abuse. *Ethics & Behavior*, 4(4), 307-329.

Fulford, J., Milton, F., Salas, D., Smith, A., Simler, A., Winlove, C., & Zeman, A. (2017). The neural correlates of visual imagery vividness—An fMRI study and literature review. *Cortex*.

Galton, F. (1880). Statistics of mental imagery. *Mind*, 5(19), 301-318.

Green, M. C., & Brock, T. C. (2002). In the mind's eye: Transportation-imagery model of narrative persuasion.

Gruenewald, T. L., Kemeny, M. E., Aziz, N., & Fahey, J. L. (2004). Acute threat to the social self: Shame, social self-esteem, and cortisol activity. *Psychosomatic medicine*, 66(6), 915-924.

Hankin, B. L. (2005). Childhood maltreatment and psychopathology: Prospective tests of attachment, cognitive vulnerability, and stress as mediating processes. *Cognitive Therapy and Research*, 29(6), 645-671.

Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007). Patients with hippocampal amnesia cannot imagine new experiences. *Proceedings of the National Academy of Sciences*, 104(5), 1726-1731.

Heim, C., & Binder, E. B. (2012). Current research trends in early life stress and depression: Review of human studies on sensitive periods, gene–environment interactions, and epigenetics. *Experimental neurology*, 233(1), 102-111.

Heim, C., Newport, D. J., Heit, S., Graham, Y. P., Wilcox, M., Bonsall, R., ... & Nemeroff, C. B. (2000). Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Jama*, 284(5), 592-597.

Hertzman, C., & Boyce, T. (2010). How experience gets under the skin to create gradients in developmental health. *Annual review of public health*, 31, 329-347.

VISUAL IMAGERY AND STRESS

Hibel, L. C., Granger, D. A., Kivlighan, K. T., Blair, C., & Family Life Project Investigators. (2006). Individual differences in salivary cortisol: Associations with common over-the-counter and prescription medication status in infants and their mothers. *Hormones and Behavior*, 50(2), 293-300.

Hill, E. E., Zack, E., Battaglini, C., Viru, M., Viru, A., & Hackney, A. C. (2008). Exercise and circulating cortisol levels: the intensity threshold effect. *Journal of endocrinological investigation*, 31(7), 587-591.

Hirsch, C. R., & Holmes, E. A. (2007). Mental imagery in anxiety disorders. *Psychiatry*, 6(4), 161-165.

Holmes, E. A., Arntz, A., & Smucker, M. R. (2007). Imagery rescripting in cognitive behaviour therapy: Images, treatment techniques and outcomes. *Journal of behavior therapy and experimental psychiatry*, 38(4), 297-305.

Holmes, E. A., Blackwell, S. E., Burnett Heyes, S., Renner, F., & Raes, F. (2016). Mental imagery in depression: Phenomenology, potential mechanisms, and treatment implications. *Annual review of clinical psychology*, 12, 249-280.

Homer, S. R., & Deeprase, C. (2017). Voluntary and Involuntary Imagery in Social Anxiety. *Behavioural and cognitive psychotherapy*, 45(3), 285-299.

Horikawa, T., Tamaki, M., Miyawaki, Y., & Kamitani, Y. (2013). Neural decoding of visual imagery during sleep. *Science*, 340(6132), 639-642.

Hornblow, A. R., & Kidson, M. A. (1976). The visual analogue scale for anxiety: a validation study. *Australian and New Zealand Journal of Psychiatry*, 10(4), 339-341.

Hyman Jr, I. E., & Pentland, J. (1996). The role of mental imagery in the creation of false childhood memories. *Journal of memory and language*, 35(2), 101-117.

Ishai, A., Haxby, J. V., & Ungerleider, L. G. (2002). Visual imagery of famous faces: effects of memory and attention revealed by fMRI. *Neuroimage*, 17(4), 1729-1741.

Iwamitsu, Y., Shimoda, K., Abe, H., Tani, T., Okawa, M., & Buck, R. (2005). The relation between negative emotional suppression and emotional distress in breast cancer diagnosis and treatment. *Health communication*, 18(3), 201-215.

Kavanagh, D. J., Andrade, J., & May, J. (2005). Imaginary relish and exquisite torture: the elaborated intrusion theory of desire. *Psychological review*, 112(2), 446.

Keenan, K. (2000). Emotion dysregulation as a risk factor for child psychopathology. *Clinical Psychology: Science and Practice*, 7(4), 418-434.

Keogh, R., & Pearson, J. (2011). Mental imagery and visual working memory. *PloS one*, 6(12), e29221.

VISUAL IMAGERY AND STRESS

Keogh, R., & Pearson, J. (2017). The blind mind: No sensory visual imagery in aphantasia. *Cortex*.

Kosslyn, S. M., Seger, C., Pani, J. R., & Hillger, L. A. (1990). When is imagery used in everyday life? A diary study. *Journal of Mental Imagery*, 14, 131-152.

Kim, H. Y. (2013). Statistical notes for clinical researchers: assessing normal distribution (2) using skewness and kurtosis. *Restorative dentistry & endodontics*, 38(1), 52-54.

Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1995). Preliminary evidence for reduced cortisol responsivity to psychological stress in women using oral contraceptive medication. *Psychoneuroendocrinology*, 20(5), 509-514.

Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The 'Trier Social Stress Test'—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, 28(1-2), 76-81.

Kosslyn, S. M., Alpert, N. M., Thompson, W. L., Maljkovic, V., Weise, S. B., Chabris, C. F., ... & Buonanno, F. S. (1993). Visual mental imagery activates topographically organized visual cortex: PET investigations. *Journal of Cognitive Neuroscience*, 5(3), 263-287.

Kosslyn, S. M., Thompson, W. L., & Ganis, G. (2006). *The case for mental imagery*. Oxford University Press.

Kounios, J., & Holcomb, P. J. (1994). Concreteness effects in semantic processing: ERP evidence supporting dual-coding theory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 20(4), 804-823.

Krakow, B., Hollifield, M., Johnston, L., Koss, M., Schrader, R., Warner, T. D., ... & Cheng, D. (2001). Imagery rehearsal therapy for chronic nightmares in sexual assault survivors with posttraumatic stress disorder: a randomized controlled trial. *Jama*, 286(5), 537-545.

Kudielka, B. M., Hellhammer, D. H., Kirschbaum, C., Harmon-Jones, E., & Winkielman, P. (2007). Ten years of research with the Trier Social Stress Test—revisited. *Social neuroscience: Integrating biological and psychological explanations of social behavior*, 56, 83.

Kuhlmann, S., Piel, M., & Wolf, O. T. (2005). Impaired memory retrieval after psychosocial stress in healthy young men. *Journal of Neuroscience*, 25(11), 2977-2982.

Leonard, B. E., & Myint, A. (2009). The psychoneuroimmunology of depression. *Human Psychopharmacology: clinical and experimental*, 24(3), 165-175.

LePort, A. K., Mattfeld, A. T., Dickinson-Anson, H., Fallon, J. H., Stark, C. E., Kruggel, F., ... & McGaugh, J. L. (2012). Behavioral and neuroanatomical

VISUAL IMAGERY AND STRESS

investigation of highly superior autobiographical memory (HSAM). *Neurobiology of learning and memory*, 98(1), 78-92.

Libby, L. K., & Eibach, R. P. (2013). The role of visual imagery in social cognition. *The Oxford handbook of social cognition*, 147-166.

Marks, D. F. (1995). New directions for mental imagery research.

MacKisack, M., Aldworth, S., Macpherson, F., Onians, J., Winlove, C., & Zeman, A. (2016). On picturing a candle: the prehistory of imagery science. *Frontiers in psychology*, 7, 515.

McEwen, B. S. (2000). Allostasis and allostatic load: implications for neuropsychopharmacology. *Neuropsychopharmacology*, 22(2), 108-124.

McEwen, B. S. (2006). Protective and damaging effects of stress mediators: central role of the brain. *Dialogues in clinical neuroscience*, 8(4), 367.

McEwen, B. S., & Seeman, T. (1999). Protective and damaging effects of mediators of stress: elaborating and testing the concepts of allostasis and allostatic load. *Annals of the New York Academy of Sciences*, 896(1), 30-47.

McEwen, B. S., & Stellar, E. (1993). Stress and the individual: mechanisms leading to disease. *Archives of internal medicine*, 153(18), 2093-2101.

McEwen, B. S. (1998). Stress, adaptation, and disease: Allostasis and allostatic load. *Annals of the New York academy of sciences*, 840(1), 33-44.

McEwen, B. S. (2005). Stressed or stressed out: what is the difference?. *Journal of Psychiatry and Neuroscience*, 30(5), 315.

McKelvie, S. J., & Demers, E. G. (1979). Individual differences in reported visual imagery and memory performance. *British Journal of Psychology*, 70(1), 51-57.

Milani, R. M., Parrott, A. C., Turner, J. J., & Fox, H. C. (2004). Gender differences in self-reported anxiety, depression, and somatization among ecstasy/MDMA polydrug users, alcohol/tobacco users, and nondrug users. *Addictive Behaviors*, 29(5), 965-971.

Miller, D. B., & O'Callaghan, J. P. (2002). Neuroendocrine aspects of the response to stress. *Metabolism-Clinical and Experimental*, 51(6), 5-10.

Moore, S. A., Zoellner, L. A., & Mollenholt, N. (2008). Are expressive suppression and cognitive reappraisal associated with stress-related symptoms?. *Behaviour research and therapy*, 46(9), 993-1000.

Morgado, P., Freitas, D., Bessa, J. M., Sousa, N., & Cerqueira, J. J. (2013). Perceived stress in obsessive-compulsive disorder is related with obsessive but not compulsive symptoms. *Frontiers in psychiatry*, 4, 21.

VISUAL IMAGERY AND STRESS

Paivio, A. (1985). Cognitive and motivational functions of imagery in human performance. *Canadian journal of applied sport sciences. Journal canadien des sciences appliquées au sport*, 10(4), 22S-28S.

Paivio, A. (1969). Mental imagery in associative learning and memory. *Psychological review*, 76(3), 241.

Palombo, D. J., Alain, C., Söderlund, H., Khuu, W., & Levine, B. (2015). Severely deficient autobiographical memory (SDAM) in healthy adults: A new mnemonic syndrome. *Neuropsychologia*, 72, 105-118.

Pearson, J., Clifford, C. W., & Tong, F. (2008). The functional impact of mental imagery on conscious perception. *Current Biology*, 18(13), 982-986.

Pearson, J., Naselaris, T., Holmes, E. A., & Kosslyn, S. M. (2015). Mental imagery: functional mechanisms and clinical applications. *Trends in cognitive sciences*, 19(10), 590-602.

Pearson, J. (2014). New directions in mental-imagery research: the binocular-rivalry technique and decoding fMRI patterns. *Current Directions in Psychological Science*, 23(3), 178-183.

Pearson, J., Rademaker, R. L., & Tong, F. (2011). Evaluating the mind's eye: the metacognition of visual imagery. *Psychological Science*, 22(12), 1535-1542.

Piccinelli, M., & Wilkinson, G. (2000). Gender differences in depression: Critical review. *The British Journal of Psychiatry*, 177(6), 486-492.

Provençal, N., & Binder, E. B. (2015). The effects of early life stress on the epigenome: from the womb to adulthood and even before. *Experimental neurology*, 268, 10-20.

Rademaker, R. L., & Pearson, J. (2012). Training visual imagery: Improvements of metacognition, but not imagery strength. *Frontiers in psychology*, 3, 224.

Richardson, J. T. (1995). Gender differences in the Vividness of Visual Imagery Questionnaire: A meta-analysis.

Roozendaal, B. (2002). Stress and memory: opposing effects of glucocorticoids on memory consolidation and memory retrieval. *Neurobiology of learning and memory*, 78(3), 578-595.

Runge, M. S., Cheung, M. W., & D'Angiulli, A. (2017). Meta-analytic comparison of trial-versus questionnaire-based vividness reportability across behavioral, cognitive and neural measurements of imagery. *Neuroscience of Consciousness*, 2017(1), nix006.

VISUAL IMAGERY AND STRESS

Saliba, F. G., Adiwardana, N. S., Uehara, E. U., Silvestre, R. N., Leite, V. V., Faleiros, F. T., & De Gobbi, J. I. (2016). Salivary cortisol levels: the importance of clown doctors to reduce stress. *Pediatric reports*, 8(1).

Sapolsky, R. M. (1996). Stress, glucocorticoids, and damage to the nervous system: the current state of confusion. *Stress*, 1(1), 1-19.

Sauro, M. D., Jorgensen, R. S., & Teal Pedlow, C. (2003). Stress, glucocorticoids, and memory: a meta-analytic review. *Stress*, 6(4), 235-245.

Schlotz, W., Yim, I. S., Zoccola, P. M., Jansen, L., & Schulz, P. (2011). The perceived stress reactivity scale: Measurement invariance, stability, and validity in three countries. *Psychological assessment*, 23(1), 80.

Schwabe, L., Joëls, M., Roozendaal, B., Wolf, O. T., & Oitzl, M. S. (2012). Stress effects on memory: an update and integration. *Neuroscience & Biobehavioral Reviews*, 36(7), 1740-1749.

Schwabe, L., & Wolf, O. T. (2010). Learning under stress impairs memory formation. *Neurobiology of learning and memory*, 93(2), 183-188.

Seeman, T. E., Singer, B. H., Rowe, J. W., Horwitz, R. I., & McEwen, B. S. (1997). Price of adaptation—allostatic load and its health consequences: MacArthur studies of successful aging. *Archives of internal medicine*, 157(19), 2259-2268.

Selye, H. (1955). Stress and disease. *The laryngoscope*, 65(7), 500-514.

Shimada, M., Takahashi, K., Ohkawa, T., Segawa, M., & Higurashi, M. (1995). Determination of salivary cortisol by ELISA and its application to the assessment of the circadian rhythm in children. *Hormone Research in Paediatrics*, 44(5), 213-217.

Sierra, M. (2009). *Depersonalization: A new look at a neglected syndrome*. Cambridge University Press.

Slee, J. A. (1980). Individual differences in visual imagery ability and the retrieval of visual appearances. *Journal of Mental Imagery*.

Sparks, G. G., Sparks, C. W., & Gray, K. (1995). Media impact on fright reactions and belief in UFOs: The potential role of mental imagery. *Communication Research*, 22(1), 3-23.

Speckens, A. E., Hackmann, A., Ehlers, A., & Cuthbert, B. (2007). Imagery special issue: Intrusive images and memories of earlier adverse events in patients with obsessive compulsive disorder. *Journal of Behavior Therapy and Experimental Psychiatry*, 38(4), 411-422.

Starkman, M. N., Schteingart, D. E., & Schork, A. M. (1981). Depressed mood and other psychiatric manifestations of Cushing's syndrome: relationship to hormone levels. *Psychosomatic medicine*, 43(1), 3-18.

VISUAL IMAGERY AND STRESS

Step toe, A., & Kivimäki, M. (2012). Stress and cardiovascular disease. *Nature Reviews Cardiology*, 9(6), 360-370.

Swann, W. B., & Miller, L. C. (1982). Why never forgetting a face matters: Visual imagery and social memory. *Journal of Personality and Social Psychology*, 43(3), 475.

Thuma, J. R., Gilders, R., Verdun, M., & Loucks, A. B. (1995). Circadian rhythm of cortisol confounds cortisol responses to exercise: implications for future research. *Journal of Applied Physiology*, 78(5), 1657-1664.

Vannucci, M., & Mazzoni, G. (2009). Individual differences in object and spatial imagery: Personality correlates. *Personality and Individual Differences*, 46(4), 402-405.

vanOyen Witvliet, C. (1997). Traumatic intrusive imagery as an emotional memory phenomenon: A review of research and explanatory information processing theories. *Clinical Psychology Review*, 17(5), 509-536.

Verma, R., Balhara, Y. P. S., & Gupta, C. S. (2011). Gender differences in stress response: Role of developmental and biological determinants. *Industrial psychiatry journal*, 20(1), 4.

Watkins, N. W. (2017). (A) phantasia and severely deficient autobiographical memory: Scientific and personal perspectives. *Cortex*.

Williams, J. M. G., Healy, H. G., & Ellis, N. C. (1999). The effect of imageability and predicability of cues in autobiographical memory. *The Quarterly Journal of Experimental Psychology Section A*, 52(3), 555-579.

Young, J. E., Klosko, J. S., & Weishaar, M. E. (2003). *Schema therapy: A practitioner's guide*. Guilford Press.

Zeman, A. Z., Della Sala, S., Torrens, L. A., Gountouna, V. E., McGonigle, D. J., & Logie, R. H. (2010). Loss of imagery phenomenology with intact visuo-spatial task performance: A case of 'blind imagination'. *Neuropsychologia*, 48(1), 145-155.

Zeman, A. Z., Dewar, M., & Della Sala, S. (2015). Lives without imagery-Congenital aphantasia.

Zeman, A., Dewar, M., & Della Sala, S. (2016). Reflections on aphantasia. *Cortex; a journal devoted to the study of the nervous system and behavior*, 74, 336-337.